

>The following few slides are from my 10 week **Attaining Optimal Health Course** 

2011 German study of about 8000 UNVACCINATED children, newborn to 19 years. Vaccinated children have 2 to 5 times more diseases and

disorders



# A similar New Zealand study. In the U.S., they are terrified to compile and publish this data



Kemp T, Pearce N, Fitzharris P, Crane J, Fergusson D, St George I, Wickens K, Beasley R.: Is infant immunization a risk factor for childhood asthma or allergy? *Epidemiology*. 1997 Nov; 8(6): 678-80.

### Journal of Translation Science Vol 3 2017

<b>Table 3.</b> Vaccination status and health outcomes – Chronic Conditions			
Chronic Disease	Vaccinated (n=405)	Unvaccinated (n=261)	Vaccinated Kids have
Allergic rhinitis			
Yes	42(10.4%)	1 (0.4%)	2,600% more
	363 (89.6%)	260 (99.6%)	
Allergies			
Yes	90(22.2%)	18(6.9%)	320% more
No	315 (77.9%)	243 (93.1%)	
ADHD			
Yes	19 <mark>(4.7%)</mark>	3 (1.0%)	470% more
No	386 (95.3%)	258 (99.0%)	
ASD			
Yes	19 <mark>(4.7%)</mark>	3 (1.0%)	470% more
No	386 (95.3%)	258 (99.0%)	
Eczema (atopic dermatitis)			
Yes	38 (9.5%)	9 (3.6%)	264% more
No	367 (90.5%)	252 (96.4%)	
Learning Disability			
Yes	23 (5.7%)	3 (1.2%)	475% more
No	382 (94.3%)	258 (98.9%)	
Neurodevelopment Disorder			
Yes	42(10.5%)	8 (3.1%)	338% more
No	313 (89.5%)	253 (96.9%)	
Any Chronic Condition			
Yes	178 <mark>(44.0%)</mark>	65 <mark>(24.9%)</mark>	176% more
No	227 (56.0%)	196 (75.1%)	

J Transl Sci, 2017 doi: 10.15761/JTS.1000186

Mumps Outbreak in Orthodox Jewish Communities in the United States New England Journal of Medicine Nov 1st 2012

### **Study Results synopsis:**

From June 28, 2009, through June 27, 2010, a total of 3502 outbreak-related cases of mumps were reported in New York City, two upstate New York counties, and one New Jersey county. Of the 1648 cases for which clinical specimens were available, 50% were laboratory-confirmed. Orthodox Jewish persons accounted for 97% of case patients. Adolescents 13 to 17 years of age (27% of all patients) and males (78% of patients in that age group) were disproportionately affected. Among case patients 13 to 17 years of age with documented vaccination status, 89% had previously received the required two doses of a mumpscontaining vaccine, and 8% had received only one dose.

#### 3 Girls Dead, Others Hospitalized After Gardasil HPV Vaccine

#### By Contributing writer, Anthony Gucciardi

Following controversy over U.S. state legislatures requiring young girls to take Gardasil, Merck's vaccine for human papillomavirus (HPV), a number of severe side effects have been observed along with the *recent* deaths of 3 young girls.

As of January **2010**, Gardasil has been linked to 49 deaths and countless side-effects, while cancer, associated with HPV, is only responsible for 1% of all cancer deaths. Why then, is it being recommended to millions worldwide?

Out of 15 million people, 49 deaths may not seem like a lot. Unfortunately, however, there are many more cases of extreme side effects from the vaccination. In fact, the amount of adverse reactions was so high that Judicial Watch, a group that claims to expose government corruption, was forced to step in to investigate.

**Between May 2009 and September 2010 alone**, Gardasil was linked to 3,589 harmful reactions and 16 deaths. Of the 3,589 adverse reactions, many were debilitating. Permanent disability was the result of 213 cases; 25 resulted in the diagnosis of Guillain-Barre Syndrome; and there were 789 other "serious" reports according to FDA documents.

"It is simply no longer possible to believe much of the clinical research that is published, or to rely on the judgment of trusted physicians or authoritative medical guidelines. I take no pleasure in this conclusion, which I reached slowly and reluctantly over my two decades as an editor of The New England Journal of Medicine" - Marcia Angell, MD. She is a physician and author, along with being the first woman to serve as editor-in-chief of The New England Journal of Medicine

Another good Vaccine Website www.vaccine-injury.info

# The 2008 FDA Statement on Gardasil reported that 73.3% of the 'healthy' girls in the clinical trials developed "New Medical Conditions"

- Blood & Lymphatic System Disorders 2.9% = 1 in 34
- Gastrointestinal Disorders 13.4% = 1 in 7
- General & Administration Site Conditions 3.8% = 1 in 33
- Immune System Disorders 2.4% =1 in 50
- Infections & Infestations 52.9% = 1 in 2
- Injury, Poisoning, & Procedural Complications 8.0% =1 in 12
- Investigations 11.8% =1 in 9
- Musculoskeletal & Connective Tissue Disorders 6.8% =1 in 14
- Nervous System Disorders 9.4% = 1 in 10
- Pregnancy &Perinatal Conditions 2.0% = 1 in 50
- Psychiatric Disorders 4.4% =1 in 22
- Renal Disorders 2.7% =1 in 37
- Reproductive & Breast Disorders 6.8 % = 1 in 14
- Respiratory, Thoracic & Mediastinal Disorders 5.5% = 1 in 18
- Skin & Subcutaneous Tissue Disorders 7.4% = 1 in 13
- Surgical Procedures = Appendectomy 10.2% = 1 in 10

Gardasil was taken off the recommended vaccine schedule in Japan

If you do not feel that the time and money to look into these references is worth it to you, then I sincerely hope you do not have, or have influence over, any current or future children.

- Watch one (or all) of the following DVDs by Dr. Sherry Tenpenny: <u>Vaccine Basics</u>, <u>Vaccines</u>: <u>What CDC</u>
  <u>Documents and Science Reveals</u>, <u>Vaccines</u>: <u>The Risks</u>, <u>The Benefits</u>, <u>The Choices</u>, <u>Vaccine Developers</u> <u>Heroes or Villains</u>. <u>MOVIES</u>: <u>VAXXED</u> & 50 Cents a Dose
- Listen to all the presentations from an entire NVIC conference. The older sets are available from AudioTapes.com
- The last one they held was in 2009 and is available with all of the audio lectures on one DVD from me for \$25.00 They stopped selling it and gave me permission to

9

## Feb. 2<sup>nd</sup> 2015 GlobeNewswire:

- Scientific evidence has demonstrated that recently vaccinated children with live virus vaccines (such as the MMR vaccine & others) can SHED the virus for weeks or months and can infect other vaccinated and unvaccinated children and adults. This can occur even though no symptoms are being displayed.
- > Numerous published studies have proven this to be true
- Parents who vaccinate their children can therefore be putting others at risk for an undisclosed number of weeks after the vaccination. (possibly over a month)
- This would indicate that playing with a recently vaccinated (less than a month ago) child has a risk that is far greater than playing with an unvaccinated child.
- >Your physician does not read this research and is a pawn of the vaccine manufacturers and government media ignorance.

### It is my firm belief, that no sane, openminded, intelligent individual, who currently thinks vaccinations are helpful:

- ➤1) Will not change their mind after examining the other side thoroughly with some of my listed references
- ▶2) Will not become an opponent to vaccination
- >3) Will not become furious about what is going on
- ▶4) Will not gain a far better understanding of the role which greed has in our healthcare and legal systems
- "What gets us into trouble is not what we don't know, it's what we know for sure that just ain't so." Josh Billings